

RESEARCH ARTICLE

10.1002/2016JG003354

Key Points:

- We assessed bulk N deposition and isotopes in urban and agricultural valleys
- N deposition and $\delta^{15}\text{N}$ of NH_4^+ were similar; $\delta^{15}\text{N}$ of NO_3^- varied seasonally with land use
- Variable particulate N contributions may influence $\delta^{15}\text{N}$ in bulk deposition

Correspondence to:

S. J. Hall,
stevenjh@iastate.edu

Citation:

Hall, S. J., E. M. Ogata, S. R. Weintraub, M. A. Baker, J. R. Ehleringer, C. I. Czimczik, and D. R. Bowling (2016), Convergence in nitrogen deposition and cryptic isotopic variation across urban and agricultural valleys in northern Utah, *J. Geophys. Res. Biogeosci.*, 121, 2340–2355, doi:10.1002/2016JG003354.

Received 26 JAN 2016

Accepted 23 AUG 2016

Accepted article online 25 AUG 2016

Published online 14 SEP 2016

Convergence in nitrogen deposition and cryptic isotopic variation across urban and agricultural valleys in northern Utah

S. J. Hall^{1,2}, E. M. Ogata³, S. R. Weintraub⁴, M. A. Baker³, J. R. Ehleringer^{1,5}, C. I. Czimczik⁶, and D. R. Bowling^{1,5}

¹Global Change and Sustainability Center, University of Utah, Salt Lake City, Utah, USA, ²Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, Iowa, USA, ³Department of Biology and the Ecology Center, Utah State University, Logan, Utah, USA, ⁴Department of Geology and Geophysics, University of Utah, Salt Lake City, Utah, USA, ⁵Department of Biology, University of Utah, Salt Lake City, Utah, USA, ⁶Department of Earth System Science, University of California, Irvine, California, USA

Abstract The extent to which atmospheric nitrogen (N) deposition reflects land use differences and biogenic versus fossil fuel reactive N sources remains unclear yet represents a critical uncertainty in ecosystem N budgets. We compared N concentrations and isotopes in precipitation-event bulk (wet + dry) deposition across nearby valleys in northern Utah with contrasting land use (highly urban versus intensive agriculture/low-density urban). We predicted greater nitrate (NO_3^-) versus ammonium (NH_4^+) and higher $\delta^{15}\text{N}$ of NO_3^- and NH_4^+ in urban valley sites. Contrary to expectations, annual N deposition ($3.5\text{--}5.1\text{ kg N ha}^{-1}\text{ yr}^{-1}$) and inorganic N concentrations were similar within and between valleys. Significant summertime decreases in $\delta^{15}\text{N}$ of NO_3^- possibly reflected increasing biogenic emissions in the agricultural valley. Organic N was a relatively minor component of deposition ($\sim 13\%$). Nearby paired wildland sites had similar bulk deposition N concentrations as the urban and agricultural sites. Weighted bulk deposition $\delta^{15}\text{N}$ was similar to natural ecosystems ($-0.6 \pm 0.7\%$). Fine atmospheric particulate matter ($\text{PM}_{2.5}$) had consistently high values of bulk $\delta^{15}\text{N}$ ($15.6 \pm 1.4\%$), $\delta^{15}\text{N}$ in NH_4^+ ($22.5 \pm 1.6\%$), and NO_3^- ($8.8 \pm 0.7\%$), consistent with equilibrium fractionation with gaseous species. The $\delta^{15}\text{N}$ in bulk deposition NH_4^+ varied by more than 40‰, and spatial variation in $\delta^{15}\text{N}$ within storms exceeded 10‰. Sporadically high values of $\delta^{15}\text{N}$ were thus consistent with increased particulate N contributions as well as potential N source variation. Despite large differences in reactive N sources, urban and agricultural landscapes are not always strongly reflected in the composition and fluxes of local N deposition—an important consideration for regional-scale ecosystem models.

1. Introduction

Atmospheric nitrogen (N) deposition derived from anthropogenic sources represents a significant N input to most regions on Earth [Galloway *et al.*, 2004]. Estimating these fluxes and ascertaining their sources over local and regional scales is critical for assessing ecosystem impacts of N as well as for informing environmental policy and regulatory compliance. However, this endeavor is complicated by the presence of numerous biogenic and/or anthropogenic sources—especially agricultural emissions from animal waste or fertilized soils, fossil fuel combustion, and industrial emissions—that have heterogeneous spatial and temporal distributions. Monitoring the chemical composition of precipitation represents a potentially effective strategy for assessing spatial and temporal variations in atmospheric reactive N sources.

In the United States, the National Atmospheric Deposition Program (NADP) uses a network of precipitation sampling sites explicitly located away from urban centers to estimate wet deposition of N at the national scale [Bigelow *et al.*, 2001]. Avoiding urban areas implicitly assumes that local urban emissions impact the chemical composition of precipitation at a given urban site. It is generally assumed that urban areas dominate oxidized N emissions, while agricultural areas dominate reduced N emissions to the atmosphere [Hertel *et al.*, 2012]. However, the impacts and relative importance of urban and agricultural emission hot spots on the spatial and temporal distributions of N deposition remain understudied.

Discrepancies between estimated reactive N emissions and measured deposition fluxes suggest that existing sampling networks have undersampled N deposition associated with urban and agricultural emission hot

spots [Holland *et al.*, 2005]. Total N deposition is typically higher in urban versus wildland areas [Fenn and Bytnerowicz, 1997; Lovett *et al.*, 2000; Fang *et al.*, 2011; Bettez and Groffman, 2013; Rao *et al.*, 2014]. Fossil fuel combustion is concentrated in urban areas [Gurney *et al.*, 2009] and represents the dominant source of nitrogen oxides (NO_x) to the atmosphere [Galloway *et al.*, 2004]. Most NO_x oxidizes to nitrate (NO_3^-) within hours [Beirle *et al.*, 2011] and is subsequently removed by wet or dry deposition over time scales of minutes to days [Hertel *et al.*, 2012]. Several studies documented increased dry deposition and foliar N uptake adjacent to roads due to vehicular NO_x emissions [Ammann *et al.*, 1999; Pearson *et al.*, 2000; Redling *et al.*, 2013]. Fertilized agricultural soils also represent a major NO_x source, particularly in semiarid climates where low moisture enhances microbial NO production [Davidson and Kinglerlee, 1997; Hertel *et al.*, 2012]. The impacts of agricultural NO_x on NO_3^- deposition have received little attention in comparison with fossil fuel combustion.

Ammonium (NH_4^+) in soluble and particulate forms, in equilibrium with gaseous NH_3 emitted to the atmosphere, represents the other primary component of bulk N deposition. Together, these species (NH_x) have typical atmospheric residence times of hours to days; particulate NH_4^+ has lower dry deposition velocities and longer residence times [Hertel *et al.*, 2012]. Agricultural regions with intensive livestock production and fertilizer application are the primary source of global NH_3 emissions [Galloway *et al.*, 2004]. However, urban areas increasingly represent a source of NH_3 from vehicular fossil fuel combustion [Kean *et al.*, 2000] in addition to smaller sources from industry and other organic wastes (e.g., landfills and sewage treatment plants) [Battye *et al.*, 2003]. Organic N may contribute an important but highly variable component to total deposition [Cape *et al.*, 2011], although relationships between organic N and land use remain poorly characterized.

Despite the strong association between land use and emissions of NO_x and NH_3 [Hertel *et al.*, 2012], relatively few studies have assessed impacts of intensive urban and agricultural land use on regional-scale (i.e., scales of tens to hundreds of kilometers) patterns of bulk N deposition [Lohse *et al.*, 2008; Tulloss and Cadenasso, 2015]. Previous studies often used forested urban to rural gradients, with natural or low-intensity agricultural landscapes as the end-member [Fenn and Bytnerowicz, 1997; Lovett *et al.*, 2000; Fang *et al.*, 2011; Bettez and Groffman, 2013; Rao *et al.*, 2014]. This work documented increased N deposition in proximity to urban centers, usually by measuring N inputs in throughfall from forested canopies. Plant canopies can more effectively scavenge atmospheric dry and fog deposition than a single planar surface (i.e., pavement or bare soil). The presence of a plant canopy typically increases deposition fluxes relative to “bulk deposition,” defined here as wet + dry deposition in the absence of overhanging plants [Fenn and Poth, 2004; Bettez and Groffman, 2013; Tulloss and Cadenasso, 2015]. Canopy throughfall deposition measurements clearly demonstrated the potential importance of urban areas as hot spots of dry N deposition.

Despite the useful information gained from throughfall deposition measurements, this method is less practical in heterogeneous urban and agricultural landscapes. First, plant species differ in their capacity to scavenge atmospheric N [Lovett, 1994; Fenn and Bytnerowicz, 1997; Sparks, 2009], which can confound assessments of land use impacts when vegetation covaries. Second, the deposition surface provided by a forest canopy is not necessarily representative of the broader landscape, especially in semiarid environments with few trees. Bulk deposition measurements provide a comparable metric for assessing regional differences in N deposition [Lewis *et al.*, 1984; Liu *et al.*, 2013], despite their underestimation of dry N deposition to canopies.

The stable isotope ($\delta^{15}\text{N}$) compositions of NH_4^+ and NO_3^- complement measurements of N concentrations, potentially providing insights into spatial and temporal variations in reactive N sources to the atmosphere. Values of $\delta^{15}\text{N}$ can reflect differences in sources and/or isotope fractionation during the production or emission of atmospheric reactive N. Atmospheric NH_3 from biogenic sources such as urine and manure, and also inorganic fertilizer, often has low $\delta^{15}\text{N}$ values (e.g., -60 – 0 ‰ [Frank *et al.*, 2004; Skinner *et al.*, 2006; Felix *et al.*, 2013, 2014]) as a consequence of kinetic and equilibrium fractionation during NH_3 volatilization. This isotopic variation arises both from $\delta^{15}\text{N}$ values of the source of mineralized NH_3 , as well as the extent of NH_3 volatilization [Hogberg, 1997]. For example, $\delta^{15}\text{N}$ values in animal waste and fertilizer vary between 2 – 25 ‰ and -2 – 2 ‰, respectively [Bateman and Kelly, 2007; Kendall *et al.*, 2007], but can yield volatilized NH_3 that is highly depleted in $\delta^{15}\text{N}$, e.g., -56 to -23 ‰ [Felix *et al.*, 2013]. In contrast to biogenic or fertilizer sources, NH_3 released or produced as a by-product of fossil fuel combustion may have relatively higher $\delta^{15}\text{N}$ values, e.g., -15 to -2 ‰ [Felix *et al.*, 2013], given that kinetic fractionation during a phase transition is less likely to be expressed. All else equal, greater values of $\delta^{15}\text{N}$ of




Figure 1. Precipitation sampling locations in (a) the geographic context of the intermountain western United States, (b) the Salt Lake Valley, and (c) the Cache Valley. The green squares represent the NADP sites, and the blue and red circles are the sites sampled for the present study. This color scheme is continued in the following figures. The light-grey shaded areas of Figures 1b and 1c represent urban land use.

NH_4^+ in atmospheric deposition likely reflect greater contributions of combustion-derived NH_3 relative to biogenic/fertilizer NH_3 .

Variation in $\delta^{15}\text{N}$ of NO_x and atmospheric NO_3^- has similarly been used to distinguish oxidized N contributions from fossil fuel and biogenic sources [Elliott et al., 2007; Walters et al., 2015]. Median values of $\delta^{15}\text{N}$ of NO_x from coal combustion (~ 10 – 20‰) are higher than median values of diesel and gasoline combustion (~ -20 – 5‰), which are both higher than median soil NO_x ($\sim -30\text{‰}$) [Li and Wang, 2008; Walters et al., 2015]. When comparing regions with similar fossil fuel NO_x sources, lower values of $\delta^{15}\text{N}$ in NO_3^- likely reflect an increasing biogenic emission contribution. In contrast to $\delta^{15}\text{N}$, values of $\delta^{18}\text{O}$ in NO_3^- may be controlled by atmospheric conditions during NO_x oxidation and provide less insight into NO_x sources [Michalski et al., 2003].

The capacity of $\delta^{15}\text{N}$ observations in bulk deposition to assess and record the influence of local emission sources on ecosystem N inputs remains largely unexplored. The origins of atmospheric NH_3 and NO_3^- are of particular interest in the intermountain western United States because of their contribution to the formation of fine atmospheric particulate matter ($\text{PM}_{2.5}$). In this region, winter cold air pools facilitate the accumulation of $\text{PM}_{2.5}$ dominated by ammonium nitrate and ammonium sulfate, which represents a major public health concern and a significant N source [Mangelson et al., 1997; Malek et al., 2006; Kelly et al., 2013; Hall et al., 2014]. The isotopic compositions of NO_3^- and NH_4^+ in $\text{PM}_{2.5}$ could provide insight into their sources, i.e., from soil and biogenic processes versus fossil fuel combustion.

Here we collected bulk deposition samples on a precipitation-event basis at sites spanning two nearby montane valleys in northern Utah, USA, with contrasting land use: the highly urban Salt Lake Valley and the predominantly agricultural Cache Valley (Figure 1). Native vegetation is characterized by a patchwork of grasses, shrubs, and bare soil; urban landscapes support a discontinuous and variable canopy of street trees in a landscape dominated by pavement, lawns, and buildings; and agricultural areas consist of seasonal or irrigated pastures and row crops and bare soil for much of the year. We asked: do bulk N deposition, N speciation, or N isotope composition reflect land use differences within and between valleys and over time? We predicted that bulk deposition in the urban Salt Lake Valley would contain more NO_3^- relative to NH_4^+ in deposition as a consequence of greater vehicular and industrial NO_x emissions in urban areas. We predicted that bulk deposition in the agricultural Cache Valley would have higher NH_4^+ concentrations due to the

prevalence of cattle feedlots and fertilized fields (which volatilize NH_3 that is deposited as NH_4^+ in acidic precipitation) and lower local fossil fuel NO_x emissions. Similarly, we predicted that $\delta^{15}\text{N}$ in NO_3^- and NH_4^+ would be higher in the Salt Lake Valley than in the Cache Valley due to increased fossil fuel versus biogenic/fertilizer sources. To provide further context, we compared bulk deposition N concentrations with wet deposition data from several regional NADP sites.

2. Materials and Methods

2.1. Study Area

The Salt Lake and Cache Valleys are characterized by a similar semiarid continental climate, with mean annual temperature of 11 and 9°C and annual precipitation of 410 and 500 mm, respectively, measured on the valley floors. Precipitation consists mostly of winter snow and spring and fall rains, punctuated by occasional summer storms. The Salt Lake Valley ($\sim 25 \times 50 \text{ km}^2$) is a major metropolitan area with a human population > one million, whereas the Cache Valley ($\sim 20 \times 60 \text{ km}^2$) is characterized by concentrated animal agriculture (cattle feedlots and dairies), pasture, and row crops and has similar populations of cattle and people (approximately 100,000 each or $>100 \text{ km}^{-2}$; Utah State University Cooperative Extension, unpublished data). We assumed that measurements of atmospheric N deposition at sites in these valleys would reflect local (within-valley) emissions as well as an unknown regional contribution. There are no other major urban areas or hot spots of concentrated feedlots/agriculture within hundreds of kilometers of these valleys. In the Salt Lake Valley, within-valley fossil fuel combustion, and dilution and mixing with incoming cleaner air masses, can explain the majority of the temporal variation in carbon dioxide (CO_2) concentrations [Strong *et al.*, 2011]. This finding implies that local (as opposed to regional) emissions might also be strongly linked to atmospheric reactive N concentrations in these valleys, especially given the shorter residence times of reactive N relative to CO_2 . Montane valleys are an especially useful spatial scale of analysis in this region during winter, where temperature inversions allow mixing of local atmospheric emissions throughout the valleys over time scales of hours to days [Pataki *et al.*, 2005].

In each valley, site locations were chosen to characterize atmospheric inputs to intensively monitored watersheds in the iUTAH hydrologic observatory (www.iUTAHepscor.org). Salt Lake Valley sites ($n = 6$) were located within or adjacent to the Red Butte Creek watershed, a focal iUTAH watershed. Five of these sites were located in urban Salt Lake City and one in Red Butte Canyon Research Natural Area, an adjacent protected area (Figure 1b). Cache Valley sites ($n = 5$) were located within the Logan River watershed, another iUTAH focal area. Three sites were located in the City of Logan (population $\sim 50,000$), one along a roadside in an agricultural landscape with pasture and row crops, and another in Logan River Canyon east of the Cache Valley. We predicted that the deposition sampled in the urban Cache Valley sites would also reflect emissions from the adjacent agricultural landscape, which stretched for 20–30 km to the south, west, and north of Logan. In both valleys, urban sampling sites were located in fenced properties to prevent tampering with equipment. Two Cache Valley urban sites were relocated to nearby locations in April 2014 due to logistical constraints.

In Salt Lake City and the City of Logan, the tree canopy consisted largely of deciduous species native to the eastern United States and represented a relatively small portion of total land cover ($\sim 26\%$) [Nowak *et al.*, 1996]. The wildland sites sampled here had similarly sparse tree cover and were dominated by low-statured ($<1 \text{ m}$) herbaceous vegetation and shrubs. The agricultural Logan Valley sites were characterized by annual row crops and pasture.

2.2. Precipitation Sampling

Snow and rain samples in the Salt Lake Valley were collected on a precipitation event basis, whenever possible, from December 2013 to February 2015 (total $n = 324$ samples). Samples were collected within 36 h of the end of precipitation events. In cases where samples could not be collected within this interval, samples were discarded and a clean collection bottle was installed to minimize the potential for microbial N immobilization. In the Cache Valley, snow and rain samples were collected from December 2013 to February 2014 and from April 2014 to February 2015 (total $n = 168$ samples). Snow was collected from acid-washed high-density polyethylene (HDPE) surfaces mounted on storm boards using acid-washed polyvinylchloride cores as described by Hall *et al.* [2014]. Rain was collected in HDPE funnels (20 cm diameter) mounted 1 m above the ground on steel posts, installed away from overhanging vegetation. Funnels were

connected via Tygon tubing to HDPE bottles placed inside plastic cylinders buried beneath the soil. Funnel necks were plugged with polyester fiber to reduce infiltration of debris and replaced following rainfall events. Samples from March 2014 often represented a mixture of snow and rain and were separately collected in acid-washed HDPE buckets (18.9 L). After collection, samples were weighed to calculate precipitation amount, filtered through precombusted Whatman GF/F filters, and frozen until analysis. Daily time series of precipitation amount were obtained from rain gauges on the University of Utah and Utah State University campuses. Values of pH measured on a subset of samples were <6 , indicating that NH_3 volatilization was minimal.

Precipitation samples represented bulk (i.e., wet + dry) deposition, given that funnel and storm board surfaces were exposed to the atmosphere between precipitation events. Our protocol contrasted with precipitation sampling at National Atmospheric Deposition Program (NADP) National Trends Network sites, where containers were exposed to the atmosphere only during precipitation events using an automated sampler. To further investigate dry deposition contributions to bulk ion loading at our sampling sites, we rinsed the funnel collectors with 18.2 M Ω deionized water for ion analysis after periods of more than 2 weeks without rain. Given that there were only three such events during the study period, we added the ion fluxes from rinsed samples to ion loads from the subsequent precipitation event for ease of interpretation.

We also compared our observations with five nearby (i.e., within several hundred of kilometers) NADP sites in the intermountain western United States (Figure 1a): Pinedale, WY; Craters of the Moon National Monument, ID; Great Basin National Park, NV; Green River, UT; and Logan, UT. The Logan NADP site was 10 km southwest of our Logan collection sites, in close proximity (<100 m) to animal feedlots. Samples from NADP sites were collected over weekly intervals according to protocols detailed in the National Trends Network operations manual (http://nadp.sws.uiuc.edu/cal/PDF/NTN_Operations_Manual_v_2-2.pdf). We analyzed the data corresponding to our period of sample collection (detailed above).

2.3. Chemical and Isotope Analyses

All samples were analyzed for NO_3^- and NH_4^+ concentrations by ion chromatography (Metrohm Compact IC, Riverview, FL). Analytical precision of ion measurements was assessed using the relative standard deviations of certified standards analyzed as unknowns. Cumulative relative standard deviations of these standards measured $<3\%$. Mean-reported values of National Institute of Standards and Technology (NIST)-traceable certified standards differed by $<4\%$ from measured values calculated using a separate set of NIST-traceable standards. Total dissolved N concentrations were measured on a subset of samples ($n = 45$) with a Shimadzu TOC-V analyzer (Columbia, MD). Dissolved organic N (DON) was calculated as the difference between total dissolved N and inorganic N ($\text{NH}_4^+ + \text{NO}_3^- + \text{NO}_2^-$; NO_2^- was typically below detection limit and not discussed further). This approach occasionally yielded negative values ($n = 6$), likely due to compounding analytical variability in the three separate chemical determinations (NH_4^+ , NO_3^- , and total dissolved N). The most negative DON sample measured -0.11 mg L^{-1} ; thus, we assigned all samples with DON absolute values $<0.11 \text{ mg L}^{-1}$ to zero to avoid positively biasing our results, as opposed to simply removing negative values.

A subset of precipitation samples ($n = 162$) was analyzed for N isotope ratios ($\delta^{15}\text{N}$) of NH_4^+ using an NH_3 diffusion method modified from Holmes *et al.* [1998]. Briefly, 30 mL of sample was added to a 60 mL HDPE bottle along with 1.5 g NaCl (to decrease the osmotic potential of the solution). A glass-fiber filter acidified with 30 μL of 4 M H_3PO_4 was sandwiched between Teflon tape and added to the bottle. Then, 90 mg of magnesium oxide was added to volatilize NH_3 , and the bottle was immediately capped and incubated for 7 days on an orbital shaker/incubator at 40°C to allow NH_3 to be completely trapped as NH_4^+ on the acidified filter. Filters were analyzed for $\delta^{15}\text{N}$ values (precision $<0.2\text{‰}$) by combustion on an elemental analyzer coupled to an isotope ratio mass spectrometer (Finnigan MAT Delta S, San Jose, CA) at the Stable Isotope Ratio Facility for Environmental Research Facility at the University of Utah (<http://sirfer.utah.edu>). To verify a lack of fractionation during recovery of NH_4^+ on filter disks, we also similarly analyzed ammonium sulfate solutions with known $\delta^{15}\text{N}$ values. We obtained equivalent $\delta^{15}\text{N}$ values (within 0.2‰) after diffusion of these standards.

Additionally, we analyzed a subset of precipitation samples ($n = 53$) for NO_3^- isotopic composition ($\delta^{15}\text{N}$ and $\delta^{18}\text{O}$) using *Pseudomonas aureofaciens* and the denitrifier method [Casciotti *et al.*, 2002] to generate N_2O for analysis on an isotope ratio mass spectrometer. We used U.S. Geological Survey (USGS) 32 and 34 reference materials (for $\delta^{15}\text{N}$) and USGS 34 and 35 (for $\delta^{18}\text{O}$) to express the data in δ notation relative to atmospheric N_2

and Vienna SMOW for N and O, respectively. Mean accuracy was 0.34‰ for $\delta^{15}\text{N}$ and 0.39‰ for $\delta^{18}\text{O}$, assessed by comparing the difference between measured versus known values of USGS 35 (for $\delta^{15}\text{N}$) and USGS 32 (for $\delta^{18}\text{O}$) analyzed as unknowns. Sample precision, determined by standard deviations of several samples analyzed in triplicate, was 0.08‰ for $\delta^{15}\text{N}$ and 0.26‰ for $\delta^{18}\text{O}$.

2.3.1. Nitrogen Isotope Ratios of $\text{PM}_{2.5}$

We sampled $\text{PM}_{2.5}$ between January and February of 2015 at sites in urban Salt Lake City and near Logan, UT (in the Cache Valley), using calibrated high-volume air samplers at Utah Department of Air Quality (DAQ) sampling stations. Samples were collected over 2–5 day intervals, for a total of six Salt Lake City samples and four Logan samples. Salt Lake City samples were collected on 20 cm \times 25 cm quartz microfiber filters (2500 QAT-UP, Pallflex Tissuquartz, Port Washington, NY). A subset of Salt Lake City samples and the Logan samples were collected on Teflon filters on DAQ-operated $\text{PM}_{2.5}$ samplers. Comparisons of $\delta^{15}\text{N}$ of NH_4^+ extracted from quartz and Teflon filters sampled in Salt Lake City (see details below) showed no significant differences. Prior to sampling, quartz filters were precombusted at 500°C for 4 h and stored in aluminum foil inside sealed plastic bags. Blank filters were co-located inside aerosol samplers but not exposed to gas flow. During $\text{PM}_{2.5}$ sampling, larger particles were removed with slotted microquartz fiber filters (TE-230-QZ, microquartz-slotted collection substrates, Tisch Environmental, Cleves, OH) on SA-230-F impactor plates (TE-230-QZ, Tisch). After collection, filters and blanks were wrapped in aluminum foil, packed into airtight plastic bags, and stored at -20°C . For isotope analysis of NO_3^- and NH_4^+ in $\text{PM}_{2.5}$, filters were immersed for 24 h in deionized water and solutions were filtered to 0.45 μm . Values of $\delta^{15}\text{N}$ and $\delta^{18}\text{O}$ of NO_3^- were determined by the denitrifier method and $\delta^{15}\text{N}$ of NH_4^+ via diffusion as described above.

Bulk $\text{PM}_{2.5}$ subsamples collected on quartz filters were also analyzed for total N content and $\delta^{15}\text{N}$ with an elemental analyzer (NA 1500 NC, Thermo Scientific, Waltham, MA) coupled to an isotope ratio mass spectrometer (DeltaPlus, Thermo Fisher Scientific (Finnigan), Waltham, MA) at the Keck Carbon Cycle Accelerator Mass Spectrometry Facility of University of California, Irvine. Data were corrected for the N mass ($1.4 \mu\text{g}/\text{cm}^2$) and $\delta^{15}\text{N}$ ratios (-6.8%) of field blanks by isotope mass balance.

2.4. Data Analysis

We compared bulk ion concentrations between the Salt Lake and Cache Valleys using mixed-effect models, where sampling sites were treated as random effects to account for temporal autocorrelations. We separately tested for differences among sites in each valley using analysis of variance (ANOVA). We also binned the data into two seasons (November–March and April–October) to allow for seasonal comparisons between valleys while minimizing excessive post hoc comparisons over finer time scales (i.e., comparing all months). The latter time period represents the predominant growing season in our study area. We did not statistically compare our samples with NADP data given that they represent two different sample types (i.e., bulk versus wet deposition) but rather report NADP data for heuristic comparisons. We assessed cumulative bulk N deposition for 2014 as the product of precipitation amount and measured N concentrations at our sites; missing values from a particular site were replaced by the means for each event in each valley. We did not have a complete record of bulk deposition chemical composition in the Cache Valley sites from mid-February to early April 2014. Total bulk deposition N concentrations were statistically equivalent between valleys during the rest of our observation period (December 2013 to February 2015). Therefore, we estimated cumulative bulk N loading for Cache Valley sites during the period of missing data as the product of precipitation amount measured at Utah State University in Logan and mean N concentrations measured during this period in the Salt Lake Valley. Gap-filled data were not used in statistical tests comparing sites or valleys.

We assessed temporal trends in precipitation ion concentrations during 2014 using generalized additive models (GAMs) with a Gaussian link function, fitted using the *mgcv* package in R [Wood, 2006]. This method allowed us to test for significant seasonal patterns and also to test whether the two valleys had significantly different temporal patterns in N concentrations or $\delta^{15}\text{N}$ —i.e., whether including a separate trend for each valley or site was statistically optimal. We fit ion concentrations as a smooth function of time using regression splines that were penalized according to their “wiggleness” during model fitting to achieve a statistically optimum degree of curvature. Curvature was expressed in terms of degrees of freedom (d.f.) analogous to the order of a polynomial, allowing for fractional d.f. The need to include differing smooth functions of time for different groups of sites (i.e., Salt Lake Valley, Cache Valley, and NADP sites) was assessed by comparing model Akaike information criterion.




Figure 2. Concentrations of (a) NO_3^- and (b) NH_4^+ in weekly precipitation samples (NADP sites; green boxes) and event-based bulk deposition samples (Salt Lake and Cache sites from this study; red and blue boxes, respectively). Sites are ordered by increasing mean NO_3^- concentrations from left to right. The boxes represent the medians and the interquartile range, and data >1.5 times the range from the box to the whiskers are denoted as outliers (circles).

the Green River NADP site ($\text{NO}_3^- = 0.34, 0.27,$ and 0.25 mg NL^{-1} and $\text{NH}_4^+ = 0.37, 0.40,$ and 0.38 mg NL^{-1} , respectively; Figure 2). The Green River site was located 250–300 km southeast of these valleys in a relatively small town (~1000 people). Together, these data suggest regional congruence of bulk deposition N inputs to these northern Utah valleys as well as the nearby landscape, which were dominated by the wet deposition component.

However, median bulk deposition N concentrations at the Salt Lake and Cache Valley sites greatly exceeded wet deposition concentrations at several other more remote NADP sites in the region (Figure 2). This contrast supports the importance of local emissions in driving N deposition to the Salt Lake and Cache Valley sites, with impacts that declined with distance. Salt Lake and Cache Valley bulk deposition inorganic N concentrations exceeded wet deposition concentrations at the Great Basin NADP site (0.16 and 0.26 mg NL^{-1} for NO_3^- and NH_4^+ , respectively; Figure 2) and were more than twofold greater than the Pinedale and Craters of the Moon NADP sites ($\text{NO}_3^- = 0.10$ and 0.08 mg NL^{-1} and $\text{NH}_4^+ = 0.12$ and 0.20 , respectively; Figure 2). In contrast to the remote NADP sites, median NO_3^- concentrations at the Logan NADP site (located in the Cache Valley) were only slightly lower (0.23 mg NL^{-1}) than the bulk deposition values measured at the nearby sites sampled in this study (0.27 mg NL^{-1}), despite the fact that they only reflected wet deposition. This finding suggests a relatively minor year-round contribution of dry deposition to the bulk deposition fluxes that we measured, which

3. Results and Discussion

3.1. Spatial and Temporal Patterns in Bulk Deposition Inorganic N Concentrations


Contrary to our hypothesis that differences in urban versus agricultural land use would affect total bulk N deposition and speciation, fluxes and concentrations of NH_4^+ and NO_3^- did not significantly differ between Salt Lake and Cache Valley sampling sites on an annual basis (Figure 2). Ammonium concentrations in bulk deposition at the Salt Lake and Cache Valley sites were 0.45 ± 0.02 and $0.51 \pm 0.04 \text{ mg NL}^{-1}$ (means and standard errors), respectively, and NO_3^- concentrations were 0.45 ± 0.03 and $0.39 \pm 0.04 \text{ mg NL}^{-1}$, respectively (Figure 2). However, bulk deposition NO_3^- concentrations (but not NH_4^+) differed seasonally between valleys. During the growing season (April–October), NO_3^- was significantly greater at the Salt Lake ($0.65 \pm 0.10 \text{ mg NL}^{-1}$) than the Cache Valley ($0.42 \pm 0.08 \text{ mg NL}^{-1}$) sites ($p < 0.01$). This pattern was predominantly driven by a peak in NO_3^- at the Salt Lake sites during June and July that was not observed at the Cache Valley sites (Figure 3).

Overall, concentrations of NH_4^+ and NO_3^- did not significantly differ among sites in a given valley, despite the inclusion of paired wildland sites (ANOVA, $p > 0.05$). Median N concentrations in bulk deposition at the Salt Lake and Cache Valley sites were also of similar magnitude to median wet deposition N concentrations at the

excluded the influence of plant canopies on dry deposition capture. Intriguingly, median Logan NADP wet deposition NH_4^+ concentrations were approximately twofold greater than our bulk deposition NH_4^+ (0.79 versus 0.40 mg N L^{-1}); possible contributing mechanisms are discussed in greater detail below.

To reiterate, we found similar annual bulk deposition N concentrations across disparate sites in the urban Salt Lake and agricultural Cache Valleys, which in turn were similar to wet deposition from the Green River NADP site. This pattern contrasts with previous work in California that documented increasing NO_3^- deposition with proximity to urban versus agricultural emission sources [Tulloss and Cadenasso, 2015]. However, our findings were similar to the results of Lohse *et al.* [2008], who found no significant differences in N deposition among wildland and urban sites upwind and downwind of Phoenix, AZ. Here overall similarities in NO_x production between the Salt Lake and Cache Valleys, indicated by similar boundary layer NO_x concentrations, may have contributed to patterns in bulk deposition NO_3^- concentrations among our study sites. Tropospheric NO_x typically has a short lifetime (hours) prior to oxidation to NO_3^- [Beirle *et al.*, 2011]. Mean NO_2 concentrations measured 14 and 11 ppb at sites in the Salt Lake and Cache Valleys, respectively, during 2014 (Utah Department of Air Quality; <http://www.airmonitoring.utah.gov/dataarchive/QL2014-NO2.pdf>). The 25th and 75th percentiles of NO_2 were 5 and 22 ppb for the

Figure 3. Inorganic N concentrations in event-based bulk deposition samples from sites in the Salt Lake Valley (blue circles; six sites) and Cache Valley (red circles; five sites) and weekly wet deposition samples from regional NADP sites (green squares; five sites). (a and b) Nitrate and NH_4^+ are shown, respectively. Samples collected in this study spanned from December 2013 to February 2015. The black lines represent the GAM fits for the Salt Lake and Cache Valley bulk deposition samples, and the grey lines represent the GAM fits for the NADP sites. (c) Daily and (d) weekly cumulative precipitations measured near our Salt Lake and Cache Valley sites are shown.



Salt Lake site and 4 and 16 ppb for the Cache Valley site. The difference in mean NO_2 concentrations between the valleys (~27% greater in Salt Lake) was very similar to the difference in median bulk deposition NO_3^- concentrations between valleys (~26% greater in Salt Lake; Figure 2), despite the fact that these differences were not statistically significant. Also, the greater peak in summertime NO_2 concentrations at the Salt Lake relative to the Cache Valley sites (indicated by the greater 75th percentile but similar 25th percentile NO_2 value) may have driven the significant summertime increase in bulk deposition NO_3^- at the Salt Lake Valley sites mentioned above.

Bulk deposition NH_4^+ concentrations also significantly increased during summer according to the temporal function fit by the GAM (Figure 3a) but did not differ between valleys on a seasonal basis. The similarity in bulk NH_4^+ concentrations between valleys was surprising given the large differences between putative urban and agricultural NH_3 sources—i.e., the prevalence of feedlots and fertilized agriculture in the Cache Valley, which are thought to dominate NH_3 emissions at the global scale [Hertel *et al.*, 2012]. A scale mismatch between the location of NH_3 emission hot spots and our measurement sites may have contributed to this result. For example, the median wet deposition NH_4^+ concentration at the Logan NADP site (0.79 mg N L^{-1}) was approximately double that of the median bulk deposition NH_4^+ concentrations at the other Cache Valley sites we measured in this study (0.40 mg N L^{-1}), despite their similar median NO_3^- concentrations. The Logan NADP site was located within 1 km of several cattle feedlots, whereas the sites that we sampled were farther away (~10 km) from this particular area.

These patterns suggest that local impacts of agricultural NH_3 emissions on wet deposition NH_4^+ may be manifested over finer spatial scales (e.g., <10 km) than reflected by valley-scale measurements. Accordingly, the sites we sampled may not have received a strong NH_3 influence from animal production despite their close proximity to other agricultural sources and the overall dominance of agricultural land use in the Cache Valley. In support of this interpretation, other studies observed declining NH_3 concentrations or bulk NH_4^+ deposition fluxes within hundreds of meters of agricultural point sources [Fowler *et al.*, 1998; Fahey *et al.*, 1999]. Previous work in the Cache Valley showed that atmospheric NH_3 concentrations varied approximately 2.5-fold between rural and urban sites separated by approximately 15 km (R. Martin *et al.*, Utah State University, unpublished data), further implying highly localized emissions. Spatial heterogeneity in atmospheric reactive N was also indicated by high spatial variation in deposition inorganic N concentrations among sites in a given valley for a given storm event. The mean within-valley standard deviations for a given sampling event (total $n = 119$) in either valley were 0.22 mg N L^{-1} for NH_4^+ and 0.13 mg N L^{-1} for NO_3^- but reached as high as 2.26 and 0.94 mg N L^{-1} , respectively. For context, mean NH_4^+ and NO_3^- concentrations measured 0.48 and 0.42 mg N L^{-1} , respectively.

Finally, in addition to patterns in emissions, the removal of atmospheric reactive N by plants [Davidson and Kingler, 1997; Sparks, 2009] may also have contributed to variability in bulk deposition values and deviation from expected land use trends. In particular, the presence of greater summer canopy cover from the row crops and pasture in the Cache Valley, compared to the smaller street-tree canopy in the Salt Lake Valley sites, may have facilitated greater net plant uptake of NH_3 and NO_2 in the former sites. This effect could have a significant impact on atmospheric N concentrations in each airshed, not just in the vicinity of the canopy, given the documented magnitude of foliar N uptake [Sparks, 2009] and a well-mixed atmospheric boundary layer. Disparities in plant canopy cover may have contributed to observed seasonal differences in NO_2 concentrations and bulk NO_3^- deposition between valleys, while partially attenuating agricultural NO_x and NH_3 emissions to the atmosphere.

As with inorganic N, annual bulk deposition DON concentrations were also similar between valleys, with means of 0.14 ± 0.04 and $0.14 \pm 0.05 \text{ mg N L}^{-1}$ in the Salt Lake and Cache Valley sites, respectively. However, 49% of the samples measured had DON concentrations below the conservative detection limit we adopted here. A global meta-analysis found that DON typically contributed between 17 and 43% (interquartile range) of total precipitation N [Cape *et al.*, 2011]. The data in our present study fell below this range (~13% of total N), lower than these previous estimates. Thus, we found no evidence that anthropogenic emissions from these valleys increased DON contributions to bulk deposition. If anything, background DON contributions may have been diluted by inorganic N from regional anthropogenic emissions.

3.2. Wet Versus Dry Contributions to Bulk N Deposition

The Salt Lake and Cache Valley bulk deposition data showed a greater proportion of extreme NO_3^- and NH_4^+ and concentration values that were ≥ 1.5 times the interquartile range (denoted in Figure 2 as circles) as compared with the NADP data, indicative of dry deposition inputs. Significant temporal differences in NO_3^- concentrations between the Salt Lake/Cache Valley bulk deposition samples and the NADP wet deposition samples were reflected by trends in the GAM smooth functions, which showed two large peaks in June/July and October/November in our bulk deposition samples relative to the NADP samples (Figure 3). These high values, especially for NO_3^- , implied the importance of increased summertime dry deposition in both valleys that was not captured by the NADP wet deposition sampling. The significant difference in summertime NO_3^- concentrations between valleys (described above) may have been driven by increased dry deposition in the Salt Lake Valley sites, which showed a pronounced NO_3^- peak in June and July. In polluted Southern California airsheds with highly seasonal precipitation inputs, dry deposition to forested canopies represents a dominant form of N inputs [Bytnerowicz and Fenn, 1996; Fenn and Bytnerowicz, 1997; Fenn and Poth, 2004]. In our study area, the temporal distribution of precipitation is more uniform (Figure 3), and significant dry NO_3^- deposition to the funnel collector surfaces was only apparent sporadically during summer and late fall.

Intriguingly, winter NO_3^- concentrations were statistically similar between the bulk deposition data obtained in this study and wet deposition data for the remote NADP sites. This suggests that little winter NO_3^- dry deposition was associated with local atmospheric reactive N emissions from the Salt Lake and Cache Valleys, despite the high concentrations of NO_3^- -rich $\text{PM}_{2.5}$ aerosols in these valleys [Mangelson et al., 1997; Kelly et al., 2013]. Previous work in the Salt Lake Valley showed that dry deposition of N to snow was highly variable in space and time [Hall et al., 2014]. Dry deposition increased during periods of persistent cold air pools, and dry deposition N inputs peaked at midelevation montane sites as opposed to the valley floor [Hall et al., 2014], where most of the sites sampled in the present study were located. Our present data provide additional evidence that winter dry N deposition may be spatially limited or sporadic in the study area.

In contrast to the temporal patterns in NO_3^- concentrations, summer NH_4^+ concentrations were similar between the Salt Lake/Cache Valley sites and the regional NADP sites (Figure 3). However, they diverged in winter, when the Salt Lake and Cache Valley samples had slightly but significantly greater NH_4^+ (note the difference between the GAM fits; Figure 3). This difference may imply a greater and more widespread influence of urban and agricultural NH_3 emissions on NH_4^+ deposition during winter, possibly due to seasonal variation in atmospheric conditions. Stable atmospheric conditions often prevail in montane valleys in the intermountain western United States during winter [Lareau et al., 2013], perhaps leading to increased atmospheric NH_x concentrations and NH_4^+ deposition relative to wildland areas.

3.3. Bulk N Deposition Totals

Cumulative bulk N deposition was similar between valleys but was variable among sites, ranging between 3.5 and 5.1 $\text{kg N ha}^{-1} \text{yr}^{-1}$ (Figure 4). Bulk deposition rates peaked in July and August, and scaled relatively closely with cumulative precipitation amount, indicating a lack of strong seasonal variation in N inputs (Figure 4). These N deposition rates are modest compared with sites impacted by urban emissions from Los Angeles, CA: $\sim 30\text{--}90 \text{ kg N ha}^{-1} \text{yr}^{-1}$, measured using throughfall deposition under forested canopies [Fenn et al., 2003]. However, cumulative N deposition in our study equaled or exceeded measurements and estimates from many remote wildland sites in the western United States ($\sim 1\text{--}4 \text{ kg N ha}^{-1} \text{yr}^{-1}$) [Williams and Tonnessen, 2000; Baron et al., 2011] and was comparable to wet + dry deposition measurements from the Phoenix, AZ, metropolitan area [Lohse et al., 2008]. The bulk N deposition totals we measured here were similar to modeled wet deposition estimates for our study region ($4\text{--}6 \text{ kg N ha}^{-1} \text{yr}^{-1}$) from 2011 to 2013, derived by combining sparse measurements with an atmospheric model [Schwede and Lear, 2014]. Our data suggest that local differences in urban versus agricultural land use do not necessarily affect these regional wet-deposition estimates. However, if N deposition was greater at our valley sites proximate to emission sources as compared with more remote montane areas in the region (i.e., tens of kilometers away [Hall et al., 2014]), our data suggest that this current model might overestimate wet deposition.

Although we saw significant evidence for summer dry deposition NO_3^- inputs as described above, our data did not reflect the very high dry deposition N inputs modeled for this region ($> 12 \text{ kg N ha}^{-1} \text{yr}^{-1}$; NADP total




Figure 4. (a) Cumulative N deposition and (b) precipitation measured during 2014 at sites in the Salt Lake Valley (blue circles; six sites) and Cache Valley (red circles; five sites).

1 mg NL⁻¹, respectively). This variability is indicated by the vertical distance among points of the same color on a given sampling date in Figure 5. This temporal variability is consistent with fine-scale spatial heterogeneity in NH₃ sources as discussed above and variable contributions of particulate N (as discussed below). It also likely reflects spatial and temporal variations in isotope fractionation during scavenging of NH_x from the atmosphere over the course of precipitation events [Heaton, 1987; Heaton et al., 1997; Xiao et al., 2012], which can function analogous to a Rayleigh distillation. Biogenic NH₃ is typically thought to have lower $\delta^{15}\text{N}$ values than fossil fuel NH₃, but few measurements have constrained this variation. Importantly, $\delta^{15}\text{N}$ values of these sources potentially overlap [Frank et al., 2004; Skinner et al., 2006; Felix et al., 2013]. This source

deposition maps; http://nadp.isws.illinois.edu/committees/tdep/tdepmaps/preview.aspx#n_wd). This discrepancy likely reflects the fact that the HDPE funnel collectors used here conservatively sampled dry deposition relative to other more complex natural surfaces but also suggests that the model may have overestimated the total area-weighted dry deposition for this region, which has highly variable canopy cover that changes with season, land use type, and landscape position (valley versus montane). Multilayered plant canopies likely would have scavenged much more dry N deposition from the atmosphere than our funnel collectors [Fenn and Poth, 2004; Sparks, 2009], but we suggest that our bulk deposition estimates provide a useful metric for assessing total deposition to the monolayered surfaces—bare soil, rock, pavement, and rooftops—that represent a substantial portion of urban, agricultural, and wildland landscapes in our semiarid study region during much of the year.

3.4. Stable Isotope Composition of Inorganic N Deposition and PM_{2.5}

Contrary to our hypothesis that differences in biogenic and fossil fuel reactive N sources would be reflected in N isotope compositions in the Salt Lake and Cache Valleys, we found no significant differences in $\delta^{15}\text{N}$ values between valleys for either bulk deposition NH₄⁺ (Figure 5a) or PM_{2.5} (discussed below). Bulk deposition $\delta^{15}\text{N}$ of NH₄⁺ varied widely, between -17.4 and 24.1‰ (mean = $-0.8 \pm 0.5\text{‰}$), but showed no consistent seasonal trend according to the GAM smooth function of $\delta^{15}\text{N}$ values over time (Figure 5a). We occasionally observed extremely high variability in $\delta^{15}\text{N}$ of NH₄⁺ and NH₄⁺ concentrations among sites within a valley during a single storm event (exceeding 10‰ and




Figure 5. Values of $\delta^{15}\text{N}$ in (a) NH_4^+ and (b and c) NO_3^- in a subset of precipitation samples from the Salt Lake Valley (blue circles) and Cache Valley (red circles).

variability, along with the spatial heterogeneity of biogenic NH_3 emissions and highly localized deposition described above, may have obscured isotopic differences between valleys despite their probable differences in NH_3 sources.

In contrast to NH_4^+ , $\delta^{15}\text{N}$ of NO_3^- differed between valleys on a seasonal basis (Figure 5b). Values of ^{15}N of NO_3^- in bulk deposition were significantly lower ($p < 0.01$) in the Cache Valley ($-4.5 \pm 0.8\text{‰}$) than in the Salt Lake Valley ($-1.3 \pm 0.5\text{‰}$) during the growing season (April–October) as assessed with ANOVA, despite the fact that a single GAM function for both valleys adequately described the overall annual trend. This summertime decrease is consistent with an increased summertime biogenic NO_x contribution from fertilized Cache Valley agricultural soils, as NO_x derived from soil microbes tends to have lower $\delta^{15}\text{N}$ values than most fossil fuel sources [Li and Wang, 2008; Walters *et al.*, 2015]. On an annual basis, $\delta^{15}\text{N}$ of NO_3^- measured $0.0 \pm 0.5\text{‰}$ and was statistically equivalent between valleys. Considering both valleys together, $\delta^{15}\text{N}$ of NO_3^- showed a significant seasonal trend (indicated by the GAM fit) with lower values during the growing season (Figure 5b). Freyer [1991] found a similar $\sim 5\text{‰}$ summertime decline in $\delta^{15}\text{N}$ of NO_3^- in European precipitation and speculated that temperature-dependent isotope exchange reactions among atmospheric NO_y species and decreased contributions of particulate NO_3^- could also be responsible for seasonal patterns in $\delta^{15}\text{N}$ of NO_3^- , in addition to potential changes in NO_x sources. Values of $\delta^{18}\text{O}$ in bulk deposition NO_3^- varied between 63.7 and 92.6‰ (mean $73.2 \pm 0.9\text{‰}$) and also displayed a significant seasonal trend of lower values during the growing season (Figure 5c). This pattern is consistent with seasonal variation in atmospheric HNO_3 formation pathways [Michalski *et al.*, 2003] and does not necessarily imply variation in NO_x sources to the atmosphere.

Mean $\delta^{15}\text{N}$ values in bulk $\text{PM}_{2.5}$ ($15.6 \pm 1.4\text{‰}$) and the NO_3^- ($8.8 \pm 0.7\text{‰}$) and NH_4^+ ($22.5 \pm 1.7\text{‰}$) components of $\text{PM}_{2.5}$ were significantly greater ($p < 0.0001$) than bulk deposition $\delta^{15}\text{N}$ of NO_3^- and NH_4^+




Figure 6. Boxplots comparing $\delta^{15}\text{N}$ values in NH_4^+ and NO_3^- of bulk deposition and $\text{PM}_{2.5}$. Bulk N of $\text{PM}_{2.5}$ represents the total N pool.

with the most enriched values of $\delta^{15}\text{N}$ of NH_4^+ in bulk deposition (Figure 6). The bulk deposition $\delta^{15}\text{N}$ values $>10\text{‰}$ exceed most previous wet deposition measurements [Heaton, 1987; Fukuzaki and Hayasaka, 2009; Jia and Chen, 2010; Xiao et al., 2012]. Concentrations of the ammonium sulfate portion of $\text{PM}_{2.5}$ remain relatively consistent throughout the year in our study area, despite the fact that ammonium nitrate and total $\text{PM}_{2.5}$ mass increase during winter [Hand et al., 2012]. Thus, one plausible interpretation of the occasionally high values of $\delta^{15}\text{N}$ of NH_4^+ observed in bulk deposition (Figure 6) was that they reflected an increasing contribution of particulate NH_4^+ (i.e., NH_4^+ derived from $\text{PM}_{2.5}$ or larger aerosols), in addition to potential variation in sources. Importantly, $\delta^{15}\text{N}$ values of bulk deposition NH_4^+ frequently exceeded 0‰ , greater than many previous estimates of $\delta^{15}\text{N}$ values for biogenic or fossil fuel NH_3 sources described above. A variable contribution of $\delta^{15}\text{N}$ -enriched particulates thus provides a plausible explanation for sporadically high $\delta^{15}\text{N}$ values of deposition NH_4^+ , an interpretation that appears to have received little attention in the literature.

This interpretation is bolstered by contrasting our results with wet deposition $\delta^{15}\text{N}$ data from China, Japan, and South Africa [Heaton, 1987; Fukuzaki and Hayasaka, 2009; Jia and Chen, 2010; Xiao et al., 2012]. These studies consistently found $\delta^{15}\text{N}$ of NH_4^+ values $<0\text{‰}$ for precipitation samples with no dry deposition component. These studies were also located in areas impacted by mixtures of urban and agricultural NH_3 sources, as were our study sites; consequently, the large positive $\delta^{15}\text{N}$ deviations we observed in some samples could be parsimoniously explained by a variable contribution of dry particulate NH_4^+ deposition collected in our bulk samplers, which was not sampled in these wet deposition studies. Similarly, values of $\delta^{15}\text{N}$ in bulk deposition NO_3^- occasionally reached as high as 8.7‰ , similar to the mean values in $\text{PM}_{2.5}$ ($8.8 \pm 0.7\text{‰}$; Figure 6). Therefore, our bulk deposition NO_3^- $\delta^{15}\text{N}$ values could also potentially be interpreted as a variable mixture of $\delta^{15}\text{N}$ -enriched particulate NO_3^- and $\delta^{15}\text{N}$ -depleted gaseous HNO_3 , in addition to a mixture of isotopically variable NO_3^- sources as discussed above.

Overall, the mass- and volume-weighted isotopic compositions and standard errors of $\delta^{15}\text{N}$ in NH_4^+ and NO_3^- in bulk deposition measured -1.6 ± 0.4 and $0.8 \pm 0.5\text{‰}$, respectively. Given that volume-weighted mean concentrations of $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$ measured 0.34 ± 0.01 and $0.24 \pm 0.01 \text{ mg NL}^{-1}$, respectively, weighted $\delta^{15}\text{N}$ of total inorganic N measured $-0.6 \pm 0.7\text{‰}$. Because relatively few comprehensive studies of the isotope composition of bulk N deposition have been conducted, these measurements provide a useful

(Figure 6). The NH_4^+ component of $\text{PM}_{2.5}$ had especially high $\delta^{15}\text{N}$ values. These measured 24.1 ± 2.3 and $20.1 \pm 1.6\text{‰}$ in the Salt Lake and Cache Valley sites, respectively, but differences were not statistically significant. Values of $\delta^{15}\text{N}$ of NO_3^- in $\text{PM}_{2.5}$ were much lower than NH_4^+ of $\text{PM}_{2.5}$ but were still significantly greater than $\delta^{15}\text{N}$ of NO_3^- observed in bulk deposition (Figure 6). In the Salt Lake and Cache Valleys, $\delta^{15}\text{N}$ of NO_3^- in $\text{PM}_{2.5}$ measured 9.5 ± 1.0 and 7.7 ± 0.5 , respectively, and did not significantly differ. Values of $\delta^{18}\text{O}$ in NO_3^- of $\text{PM}_{2.5}$ measured $81.6 \pm 2.7\text{‰}$, similar to bulk deposition. Annual mean concentrations of $\text{PM}_{2.5}$ were similar at monitoring sites in both study valleys during 2013 and 2014 (Utah Department of Air Quality; <http://www.airmonitoring.utah.gov/dataarchive/>).

The high $\delta^{15}\text{N}$ values of $\text{PM}_{2.5}$ are consistent with an $\sim 33\text{‰}$ equilibrium fractionation between gaseous NH_3 and particulate NH_4^+ and $\sim 21\text{‰}$ for ammonium nitrate relative to nitric acid vapor [Heaton et al., 1997]. These fractionations lead to much greater $\delta^{15}\text{N}$ in $\text{PM}_{2.5}$ relative to gaseous precursors. Individual $\text{PM}_{2.5}$ samples varied between 17.3 and 31.4‰ in $\delta^{15}\text{N}$ of NH_4^+ and overlapped

constraint for ecosystem-scale N isotope models. For example, N isotope mass balance has been used to estimate the importance of denitrification at the global scale, requiring an estimate of the N isotope composition of atmospheric deposition inputs [Houlton and Bai, 2009]. It is uncertain whether concentrated urban or agricultural N emissions might alter the isotope composition of deposition measured in more remote ecosystems. Previous studies in natural ecosystems had a median $\delta^{15}\text{N}$ value of -1.3‰ [Houlton and Bai, 2009], similar to our present data. These similarities imply that local urban or agricultural influences do not necessarily impart a significant isotopic signature to bulk deposition N inputs on an annual basis.

4. Conclusions

Despite the strong impacts of land use on NO_x and NH_3 emissions documented elsewhere [Galloway et al., 2004; Hertel et al., 2012], we conclude that substantial valley-scale differences in urban versus agricultural land use in northern Utah were not strongly manifested in the mean speciation, isotope composition, or fluxes of bulk atmospheric N deposition to our funnel collectors on an annual basis. Differences between valleys were only apparent during summer, when bulk deposition NO_3^- concentrations increased at the Salt Lake Valley sites, and $\delta^{15}\text{N}$ of NO_3^- was lower at the Cache Valley sites. Our data suggest that relatively coarse regional estimates of these parameters may suffice for use in ecosystem-scale N cycling models, although they may miss hot spots of NH_3 emissions that are deposited very close to the source. Although median-measured bulk deposition N concentrations exceeded wet deposition concentrations from several remote NADP sites in the region, they were similar to wet deposition from another regional NADP site. It is well known that plant canopies scavenge gaseous and particulate reactive N to a greater extent than the funnels we used to collect bulk deposition [Fenn and Poth, 2004; Sparks, 2009; Bettez and Groffman, 2013]. However, our bulk deposition estimates likely provide a useful metric for estimating total deposition to nonvegetated surfaces—soil, rocks, pavement, and rooftops—that represent a substantial portion of wildland, urban, and agricultural landscapes in the intermountain western United States, especially during fall and winter.

Values of $\delta^{15}\text{N}$ in atmospheric deposition are complex to interpret because they not only represent a mixture of reactive N sources varying in $\delta^{15}\text{N}$ but also a variable mixture of phases (gas, particulate, and aqueous) impacted by large isotope fractionations. Our data highlight the potential importance of the latter phenomenon. We found large differences between $\delta^{15}\text{N}$ of bulk deposition and $\text{PM}_{2.5}$, suggesting that temporal variation in particulate contributions to bulk N deposition could potentially obscure initial isotope differences among biological versus fossil fuel reactive N sources. Even where source $\delta^{15}\text{N}$ values differ, spatial and temporal heterogeneities in particulate contributions could decrease the power to detect these differences. Thus, we recommend caution when interpreting atmospheric $\delta^{15}\text{N}$ measurements solely in terms of mixtures of sources. This is especially important when sampling schemes may selectively capture gaseous versus particulate N as opposed to the entire atmospheric reactive N pool, thus leading to physical isotope fractionation.

Acknowledgments

Upon manuscript acceptance all data are publicly available online at the Hydroshare database: <http://dx.doi.org/10.4211/hs.7ce47150ee344-d4e95f2fd7bb4f660ac>. We thank four anonymous reviewers for their thoughtful comments and the NADP program for providing their data online. S. Chakraborty and D. Fernandez provided the analytical assistance and D. Eiriksson, E. Schulze, D. Tarboton, B. Greene, D. Epstein, S. Jackson, J. Turner, A. Armstrong, and H. Quinn assisted with sample collection. J. Wessling, D. Zhang, and X. Xu contributed to sample analysis. We also thank Utah Department of Air Quality staff, especially S. Arens, for assistance with $\text{PM}_{2.5}$ collection and B. Schichtel from the U.S. National Park Service for use of a high-volume sampler. This research was supported by National Science Foundation (NSF) EPSCoR grant IIA 1208732 awarded to Utah State University as part of the State of Utah Research Infrastructure Improvement Award and by NSF grant DBI-1337947 to the University of Utah. Any opinions, findings, and conclusions or recommendations expressed are those of the author (s) and do not necessarily reflect the views of the National Science Foundation.

References

- Ammann, M., R. Siegwolf, F. Pichlmayer, M. Suter, M. Saurer, and C. Brunold (1999), Estimating the uptake of traffic-derived NO_2 from ^{15}N abundance in Norway spruce needles, *Oecologia*, 118(2), 124–131, doi:10.1007/s004420050710.
- Baron, J. S., C. T. Driscoll, J. L. Stoddard, and E. E. Richer (2011), Empirical critical loads of atmospheric nitrogen deposition for nutrient enrichment and acidification of sensitive US Lakes, *BioScience*, 61(8), 602–613, doi:10.1525/bio.2011.61.8.6.
- Bateman, A. S., and S. D. Kelly (2007), Fertilizer nitrogen isotope signatures, *Isotopes Environ. Health Stud.*, 43(3), 237–247, doi:10.1080/10256010701550732.
- Battye, W., V. P. Aneja, and P. A. Roelle (2003), Evaluation and improvement of ammonia emissions inventories, *Atmos. Environ.*, 37(27), 3873–3883, doi:10.1016/S1352-2310(03)00343-1.
- Beirle, S., K. F. Boersma, U. Platt, M. G. Lawrence, and T. Wagner (2011), Megacity emissions and lifetimes of nitrogen oxides probed from space, *Science*, 333(6050), 1737–1739, doi:10.1126/science.1207824.
- Bettez, N. D., and P. M. Groffman (2013), Nitrogen deposition in and near an urban ecosystem, *Environ. Sci. Technol.*, 47(11), 6047–6051, doi:10.1021/es400664b.
- Bigelow, S. D., S. R. Dossett, and V. C. Bowersox (2001), *Instruction Manual NADP/NTN Site Selection and Installation*, National Atmospheric Deposition Program, Champaign, Ill.
- Bytnerowicz, A., and M. E. Fenn (1996), Nitrogen deposition in California forests: A review, *Environ. Pollut.*, 92(2), 127–146.
- Cape, J. N., S. E. Cornell, T. D. Jickells, and E. Nemitz (2011), Organic nitrogen in the atmosphere—Where does it come from? A review of sources and methods, *Atmospheric Res.*, 102(1–2), 30–48, doi:10.1016/j.atmosres.2011.07.009.
- Casciotti, K. L., D. M. Sigman, M. G. Hastings, J. K. Böhlke, and A. Hilkert (2002), Measurement of the oxygen isotopic composition of nitrate in seawater and freshwater using the denitrifier method, *Anal. Chem.*, 74(19), 4905–4912, doi:10.1021/ac020113w.
- Davidson, E. A., and W. Kinglerlee (1997), A global inventory of nitric oxide emissions from soils, *Nutr. Cycl. Agroecosystems*, 48(1–2), 37–50, doi:10.1023/A:1009738715891.

- Elliott, E. M., C. Kendall, S. D. Wankel, D. A. Burns, E. W. Boyer, K. Harlin, D. J. Bain, and T. J. Butler (2007), Nitrogen isotopes as indicators of NO_x source contributions to atmospheric nitrate deposition across the midwestern and northeastern United States, *Environ. Sci. Technol.*, *41*(22), 7661–7667, doi:10.1021/es070898t.
- Fahey, T. J., C. J. Williams, J. N. Rooney-Varga, C. C. Cleveland, K. M. Postek, S. D. Smith, and D. R. Bouldin (1999), Nitrogen deposition in and around an intensive agricultural district in central New York, *J. Environ. Qual.*, *28*(5), 1585, doi:10.2134/jeq1999.00472425002800050025x.
- Fang, Y., M. Yoh, K. Koba, W. Zhu, Y. Takebayashi, Y. Xiao, C. Lei, J. Mo, W. Zhang, and X. Lu (2011), Nitrogen deposition and forest nitrogen cycling along an urban–rural transect in southern China, *Glob. Change Biol.*, *17*(2), 872–885, doi:10.1111/j.1365-2486.2010.02283.x.
- Felix, J. D., E. M. Elliott, T. J. Gish, L. L. McConnell, and S. L. Shaw (2013), Characterizing the isotopic composition of atmospheric ammonia emission sources using passive samplers and a combined oxidation-bacterial denitrifier approach, *Rapid Commun. Mass Spectrom.*, *27*(20), 2239–2246, doi:10.1002/rcm.6679.
- Felix, J. D., E. M. Elliott, T. Gish, R. Maghirang, L. Cambal, and J. Clougherty (2014), Examining the transport of ammonia emissions across landscapes using nitrogen isotope ratios, *Atmos. Environ.*, *95*, 563–570, doi:10.1016/j.atmosenv.2014.06.061.
- Fenn, M. E., and A. Bytnerowicz (1997), Summer throughfall and winter deposition in the San Bernardino Mountains in Southern California, *Atmos. Environ.*, *31*(5), 673–683.
- Fenn, M. E., and M. A. Poth (2004), Monitoring nitrogen deposition in throughfall using ion exchange resin columns: A field test in the San Bernardino Mountains, *J. Environ. Qual.*, *33*(6), 2007–2014.
- Fenn, M. E., et al. (2003), Nitrogen emissions, deposition, and monitoring in the western United States, *BioScience*, *53*(4), 391–403, doi:10.1641/0006-3568(2003)053[0391:NEDAM]2.0.CO;2.
- Fowler, D., C. E. R. Pitcairn, M. A. Sutton, C. Flechard, B. Loubet, M. Coyle, and R. C. Munro (1998), The mass budget of atmospheric ammonia in woodland within 1 km of livestock buildings, *Environ. Pollut.*, *102*(1), Suppl. 1, 343–348, doi:10.1016/S0269-7491(98)80053-5.
- Frank, D. A., R. D. Evans, and B. F. Tracy (2004), The role of ammonia volatilization in controlling the natural ¹⁵N abundance of a grazed grassland, *Biogeochemistry*, *68*(2), 169–178, doi:10.1023/B:BI0G.0000025736.19381.91.
- Freyer, H. D. (1991), Seasonal variation of ¹⁵N/¹⁴N ratios in atmospheric nitrate species, *Tellus B*, *43*(1), 30–44, doi:10.1034/j.1600-0889.1991.00003.x.
- Fukuzaki, N., and H. Hayasaka (2009), Seasonal variations of nitrogen isotopic ratios of ammonium and nitrate in precipitations collected in the Yahiko–Kakuda Mountains area in Niigata Prefecture, Japan, *Water Air Soil Pollut.*, *203*(1–4), 391–397, doi:10.1007/s11270-009-0026-8.
- Galloway, J. N., et al. (2004), Nitrogen cycles: Past, present, and future, *Biogeochemistry*, *70*(2), 153–226, doi:10.1007/s10533-004-0370-0.
- Gurney, K. R., D. L. Mendoza, Y. Zhou, M. L. Fischer, C. C. Miller, S. Geethakumar, and S. de la Rue du Can (2009), High resolution fossil fuel combustion CO₂ emission fluxes for the United States, *Environ. Sci. Technol.*, *43*(14), 5535–5541, doi:10.1021/es900806c.
- Hall, S. J., G. Maurer, S. W. Hoch, R. Taylor, and D. R. Bowling (2014), Impacts of anthropogenic emissions and cold air pools on urban to montane gradients of snowpack ion concentrations in the Wasatch Mountains, Utah, *Atmos. Environ.*, *98*, 231–241, doi:10.1016/j.atmosenv.2014.08.076.
- Hand, J. L., B. A. Schichtel, M. Pitchford, W. C. Malm, and N. H. Frank (2012), Seasonal composition of remote and urban fine particulate matter in the United States, *J. Geophys. Res.*, *117*, D05209, doi:10.1029/2011JD017122.
- Heaton, T. H. E. (1987), ¹⁵N/¹⁴N ratios of nitrate and ammonium in rain at Pretoria, South Africa, *Atmospheric Environ.* *1967*, *21*(4), 843–852, doi:10.1016/0004-6981(87)90080-1.
- Heaton, T. H. E., B. Spiro, and S. M. C. Robertson (1997), Potential canopy influences on the isotopic composition of nitrogen and sulphur in atmospheric deposition, *Oecologia*, *109*(4), 600–607, doi:10.1007/s004420050122.
- Hertel, O., et al. (2012), Governing processes for reactive nitrogen compounds in the European atmosphere, *Biogeosciences*, *9*(12), 4921–4954, doi:10.5194/bg-9-4921-2012.
- Hogberg, P. (1997), Tansley review no. 95. ¹⁵N natural abundance in soil-plant systems, *New Phytol.*, *137*, 179–203.
- Holland, E. A., B. H. Braswell, J. Sulzman, and J.-F. Lamarque (2005), Nitrogen deposition onto the United States and Western Europe: Synthesis of observations and models, *Ecol. Appl.*, *15*(1), 38–57, doi:10.1890/03-5162.
- Holmes, R. M., J. W. McClelland, D. M. Sigman, B. Fry, and B. J. Peterson (1998), Measuring ¹⁵N–NH₄⁺ in marine, estuarine and fresh waters: An adaptation of the ammonia diffusion method for samples with low ammonium concentrations, *Mar. Chem.*, *60*(3–4), 235–243, doi:10.1016/S0304-4203(97)00099-6.
- Houlton, B. Z., and E. Bai (2009), Imprint of denitrifying bacteria on the global terrestrial biosphere, *Proc. Natl. Acad. Sci. U.S.A.*, *106*(51), 21,713–21,716, doi:10.1073/pnas.0912111106.
- Jia, G., and F. Chen (2010), Monthly variations in nitrogen isotopes of ammonium and nitrate in wet deposition at Guangzhou, south China, *Atmos. Environ.*, *44*(19), 2309–2315, doi:10.1016/j.atmosenv.2010.03.041.
- Kean, A. J., R. A. Harley, D. Littlejohn, and G. R. Kendall (2000), On-road measurement of ammonia and other motor vehicle exhaust emissions, *Environ. Sci. Technol.*, *34*(17), 3535–3539, doi:10.1021/es991451q.
- Kelly, K. E., R. Kotchenruther, R. Kuprov, and G. D. Silcox (2013), Receptor model source attributions for Utah's Salt Lake City airshed and the impacts of wintertime secondary ammonium nitrate and ammonium chloride aerosol, *J. Air Waste Manag. Assoc.*, *63*(5), 575–590, doi:10.1080/10962247.2013.774819.
- Kendall, C., E. M. Elliott, and S. D. Wankel (2007), Tracing anthropogenic inputs of nitrogen to ecosystems, in *Stable Isotopes in Ecology and Environmental Science*, edited by R. Michener and K. Lajtha, pp. 375–449, Blackwell Ltd, N. J.
- Lareau, N. P., E. Crosman, C. D. Whiteman, J. D. Horel, S. W. Hoch, W. O. J. Brown, and T. W. Horst (2013), The persistent cold-air pool study, *Bull. Am. Meteorol. Soc.*, *94*(1), 51–63, doi:10.1175/BAMS-D-11-00255.1.
- Lewis, W. M., M. C. Grant, and J. F. Saunders (1984), Chemical patterns of bulk atmospheric deposition in the state of Colorado, *Water Resour. Res.*, *20*(11), 1691–1704, doi:10.1029/WR020i01p01691.
- Li, D., and X. Wang (2008), Nitrogen isotopic signature of soil-released nitric oxide (NO) after fertilizer application, *Atmos. Environ.*, *42*(19), 4747–4754, doi:10.1016/j.atmosenv.2008.01.042.
- Liu, X., et al. (2013), Enhanced nitrogen deposition over China, *Nature*, *494*(7438), 459–462, doi:10.1038/nature11917.
- Lohse, K. A., D. Hope, R. Sponseller, J. O. Allen, and N. B. Grimm (2008), Atmospheric deposition of carbon and nutrients across an arid metropolitan area, *Sci. Total Environ.*, *402*(1), 95–105, doi:10.1016/j.scitotenv.2008.04.044.
- Lovett, G. M. (1994), Atmospheric deposition of nutrients and pollutants in North America: An ecological perspective, *Ecol. Appl.*, *4*(4), 629–650, doi:10.2307/1941997.
- Lovett, G. M., M. M. Traynor, R. V. Pouyat, M. M. Carreiro, W.-X. Zhu, and J. W. Baxter (2000), Atmospheric deposition to oak forests along an urban–rural gradient, *Environ. Sci. Technol.*, *34*(20), 4294–4300, doi:10.1021/es001077q.
- Malek, E., T. Davis, R. S. Martin, and P. J. Silva (2006), Meteorological and environmental aspects of one of the worst national air pollution episodes (January, 2004) in Logan, Cache Valley, Utah, USA, *Atmospheric Res.*, *79*(2), 108–122, doi:10.1016/j.atmosres.2005.05.003.

- Mangelson, N. F., L. Lewis, J. M. Joseph, W. Cui, J. Machir, D. J. Eatough, L. B. Rees, T. Wilkerson, and D. T. Jensen (1997), The contribution of sulfate and nitrate to atmospheric fine particles during winter inversion fogs in Cache Valley, Utah, *J. Air Waste Manag. Assoc.*, *47*(2), 167–175, doi:10.1080/10473289.1997.10464429.
- Michalski, G., Z. Scott, M. Kabilig, and M. H. Thiemens (2003), First measurements and modeling of $\Delta^{17}\text{O}$ in atmospheric nitrate, *Geophys. Res. Lett.*, *30*(16), doi:10.1029/2003GL017015.
- Nowak, D. J., R. A. Rowntree, E. G. McPherson, S. M. Sisinni, E. R. Kerkmann, and J. C. Stevens (1996), Measuring and analyzing urban tree cover, *Landsc. Urban Plan.*, *36*(1), 49–57, doi:10.1016/S0169-2046(96)00324-6.
- Pataki, D. E., B. J. Tyler, R. E. Peterson, A. P. Nair, W. J. Steenburgh, and E. R. Pardyjak (2005), Can carbon dioxide be used as a tracer of urban atmospheric transport?, *J. Geophys. Res.*, *110*, D15102, doi:10.1029/2004JD005723.
- Pearson, J., D. M. Wells, K. J. Seller, A. Bennett, A. Soares, J. Woodall, and M. J. Ingrouille (2000), Traffic exposure increases natural $\delta^{15}\text{N}$ and heavy metal concentrations in mosses, *New Phytol.*, *147*(2), 317–326, doi:10.1046/j.1469-8137.2000.00702.x.
- Rao, P., L. R. Hutyla, S. M. Raciti, and P. H. Templer (2014), Atmospheric nitrogen inputs and losses along an urbanization gradient from Boston to Harvard forest, MA, *Biogeochemistry*, *121*(1), 229–245, doi:10.1007/s10533-013-9861-1.
- Redling, K., E. Elliott, D. Bain, and J. Sherwell (2013), Highway contributions to reactive nitrogen deposition: Tracing the fate of vehicular NO_x using stable isotopes and plant biomonitors, *Biogeochemistry*, *116*(1–3), 261–274, doi:10.1007/s10533-013-9857-x.
- Schwede, D. B., and G. G. Lear (2014), A novel hybrid approach for estimating total deposition in the United States, *Atmos. Environ.*, *92*, 207–220, doi:10.1016/j.atmosenv.2014.04.008.
- Skinner, R., P. Ineson, H. Jones, D. Sleep, and M. Theobald (2006), Sampling systems for isotope-ratio mass spectrometry of atmospheric ammonia, *Rapid Commun. Mass Spectrom.*, *20*(2), 81–88, doi:10.1002/rcm.2279.
- Sparks, J. P. (2009), Ecological ramifications of the direct foliar uptake of nitrogen, *Oecologia*, *159*(1), 1–13, doi:10.1007/s00442-008-1188-6.
- Strong, C., C. Stwertka, D. R. Bowling, B. B. Stephens, and J. R. Ehleringer (2011), Urban carbon dioxide cycles within the Salt Lake Valley: A multiple-box model validated by observations, *J. Geophys. Res.*, *116*, D15307, doi:10.1029/2011JD015693.
- Tulloss, E. M., and M. L. Cadenasso (2015), Nitrogen deposition across scales: Hotspots and gradients in a California savanna landscape, *Ecosphere*, *6*(9), art167, doi:10.1890/ES14-00440.1.
- Walters, W. W., B. D. Tharp, H. Fang, B. J. Kozak, and G. Michalski (2015), Nitrogen isotope composition of thermally produced NO_x from various fossil-fuel combustion sources, *Environ. Sci. Technol.*, *49*(19), 11,363–11,371, doi:10.1021/acs.est.5b02769.
- Williams, M. W., and K. A. Tonnesen (2000), Critical loads for inorganic N deposition in the Colorado Front Range, USA, *Ecol. Appl.*, *10*(6), 1648–1665, doi:10.1890/1051-0761(2000)010[1648:CLFIND]2.0.CO;2.
- Wood, S. (2006), *Generalized Additive Models: An Introduction with R*, CRC Press, Boca Raton, Fla.
- Xiao, H.-W., H.-Y. Xiao, A. Long, and Y.-L. Wang (2012), Who controls the monthly variations of NH_4^+ nitrogen isotope composition in precipitation?, *Atmos. Environ.*, *54*, 201–206, doi:10.1016/j.atmosenv.2012.02.035.